

University of Groningen

Projector 2

van Hijum, SAFT; Zomer, AL; Kuipers, OP; Kok, J

Published in:
Nucleic Acids Res.

DOI:
[10.1093/nar/gki356](https://doi.org/10.1093/nar/gki356)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2005

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van Hijum, SAFT., Zomer, AL., Kuipers, OP., & Kok, J. (2005). Projector 2: contig mapping for efficient gap-closure of prokaryotic genome sequence assemblies. *Nucleic Acids Res.*, 33(Web Server issue), W560-W566. <https://doi.org/10.1093/nar/gki356>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Contigs successfully uploaded.

[Help, all these settings, what should I change?](#)

Processing settings

Size of chopped fragments	<input type="text" value="300"/>	(Required; integer)
Minimum size of contig (otherwise it will be discarded)	<input type="text" value="1200"/>	(Required; integer)
Minimum size of remaining fragment to use for positioning	<input type="text" value="500"/>	(Required; integer)
Cutoff e-value for repeats (is also dependent on chop size)	<input type="text" value="1e-20"/>	(Required; float)

Mapping settings

Maximum size deviation (%) of mapped vs original contig	<input type="text" value="40"/>	(Required; integer)
---	---------------------------------	-----------------------

Primer design settings

Minimum size of template for primer design	<input type="text" value="300"/>	(Required; integer)
Spacing from the end of a contig (omit bad sequence area)	<input type="text" value="150"/>	(Required; integer)